


**ROLE OF NIGELLA SATIVA IN
OPIOID DEPENDENCE**

BY

DR. SIBGHATULLAH SANGI

M.B.,B.S., M.Phil



Sibghatullah Sangi
D. Phil. Degree Class
1994

UNIVERSITY OF KARACHI

2004

Dedicated
to

MY TEACHERS

Prof. Dr. Shakida P. Ahmed,
Dr. Kausar Amir

AND MY CHILDREN

Bilal Badshah, Hadiya Shahxadi, and
Huda Rani

ACKNOWLEDGMENT

All praises be to Almighty Allah the Omnipotent, the Omnipresent whose kindness and help made me able to complete this work and made hardships coming in my way, easy to be ignored. Words are helpless to express thanks and gratitude to Allah

I feel great privilege and pleasure in acknowledging my deepest sense of gratitude to the most inspiring supervisor and a great scholar of Pharmacology Prof. Dr. Shahida P. Ahmed, Professor and Chairperson, Department of Pharmacology, for her encouragement, generous supervision, and valuable suggestions during the course of present study.

I am also thankful to Dr. Kausar Aamir, Chairperson, Department of Pharmacology and Therapeutics, BMSI, JPMC, Karachi, for her kind cooperation in all odds and evens. Without her guidance it would have been difficult for me to start and complete this task.

I also wish to acknowledge for the mutual cooperation of other faculty staff of the Pharmacology department Prof. Dr. Asif Bin Rehman, Dr. Rafeeq Alam Khan, and Raheela Najam to pursue this work on an excellent way.

I feel pleasure to thank my friends and family members for their moral help and encouragement especially Dr. Shah Murad Mastoi and Dr. Poonam Chand, Incharge Rehabilitation Centre, Malir, Muhammad Arshad, ASI Police, and Mr. M. Haroon of the Pharmacology department, University of Karachi, for their valuable support.

CONTENTS

S. No.	CONTENTS	PAGE
1.	SUMMARY (URDU)	1
2.	SUMMARY (ENGLISH)	2
3.	INTRODUCTION	4
4.	MATERIALS & METHODS	48
5.	OBSERVATIONS & RESULTS	59
6.	DISCUSSION	107
7.	REFERENCES	114
8.	APPENDICES	I – XXVIII

LIST OF TABLES

	PAGE
1. Effects of Nigella Sativa 500 mg treatment on subjective symptoms of withdrawal from opioids in Group-II patients during 12 days stay in hospital	76
2. Effects of Nigella Sativa 500 mg on objective signs of withdrawal from opioids in Group-II patients during 12 days stay in hospital	79
2A. Comparison of abstinence of heroin in Group-II patients treated with Nigella Sativa 500 mg day-3 versus day-12 scores of symptoms and signs and physiological parameters	82
3. Effects of Nigella Sativa treatment on subjective symptoms of protracted abstinence in Group-II patients with opioid dependence during 12 weeks of follow up	84
4. Effects of Nigella Sativa treatment on objective signs of protracted abstinence in Group-II patients with opioid dependence during 12 weeks of follow up	87
4A. Comparison of protracted abstinence of heroin dependence in Group-II patients treated with Nigella Sativa 500 mg day-3 versus week-12 scores of symptoms and signs and physiological parameters	90
5. Effects of Nigella Sativa treatment on subjective symptoms of withdrawal from opioids in Group-I patients during 12 days stay in hospital	61
6. Effects of Nigella Sativa 250 mg treatment on objective signs of withdrawal from opioids in Group-I patients during 12 days stay in hospital	64
6A. Comparison of abstinence of heroin in Group-I patients treated with Nigella Sativa 250 mg day-3 versus day-12 scores of symptoms and signs and physiological parameters	67
7. Effects of Nigella Sativa 250 mg treatment on subjective symptoms of protracted abstinence in Group-I patients with opioid dependence during 12 weeks of follow up	69

(Continued.....)

8.	Effects of Nigella Sativa 250 mg treatment on objective signs of protracted abstinence in Group-I patients with opioid dependence during 12 weeks of follow up	71
8A.	Comparison of protracted abstinence of heroin dependence in Group-I patients treated with Nigella Sativa 250 mg day-3 versus week-12 scores of symptoms and signs and physiological parameters	74
9.	Comparison of effects of treatment with Nigella Sativa 250 mg and 500 mg on subjective symptoms of withdrawal from opioids during 12 days' stay in hospital	93
10.	Comparison of effects of treatment with Nigella Sativa 250 mg and 500 mg on objective signs of withdrawal from opioids during 12 days stay in hospital	96
10A	Comparison of abstinence of heroin in patients treated with Nigella Sativa 250 mg and 500 mg on day-3 scores of symptoms and signs and physiological parameters	99
10B	Comparison of abstinence of heroin in patients treated with Nigella Sativa 250 mg and 500 mg on day-12 scores of symptoms and signs and physiological parameters	100
11.	Comparison of effects of treatment with Nigella Sativa 250 mg and 500 mg on subjective symptoms of protracted abstinence in patients with opioid dependence during 12 weeks of follow up	102
12.	Comparison of effects of treatment with Nigella Sativa 250 mg and 500 mg on objective signs of protracted abstinence with opioid dependence during 12 weeks of follow up	104
12A	Comparison of protracted abstinence of heroin dependence in patients treated with Nigella Sativa 250 mg and 500 mg on week-12 scores of symptoms and signs and physiological parameters	106

LIST OF FIGURES

	PAGE
1. Effects of Nigella Sativa 250 mg treatment on subjective symptoms of withdrawal from opioids in Group-I patients during 12 days stay in hospital	62
2. Effects of Nigella Sativa 250 mg treatment on objective signs of withdrawal from opioids in Group-I patients during 12 days stay in hospital	65
3. Effects of Nigella Sativa 250 mg treatment on subjective symptoms of protracted abstinence in Group-I patients with opioid dependence during 12 weeks of Follow up	70
4. Effects of Nigella Sativa 250 mg treatment on objective signs of protracted abstinence in Group-I patients with opioid dependence during 12 weeks of follow up	72
5. Effects of Nigella Sativa 500 mg treatment on subjective symptoms of withdrawal from opioids in Group-II during 12 days stay in hospital	77
6. Effects of Nigella Sativa 500 mg treatment on objective signs of withdrawal from opioids in Group-II patients during 12 days stay in hospital	80
7. Effects of Nigella Sativa 500 mg treatment on subjective symptoms of protracted abstinence in Group-II patients with opioid dependence during 12 weeks of follow up	85
8. Effects of Nigella Sativa 500 mg treatment on objective signs of protracted abstinence in Group-II patients with opioid dependence during 12 weeks of follow up	88
9. Comparison of effects of treatment with Nigella Sativa 250 mg and 500 mg on subjective symptoms of withdrawal from opioids during 12 days' stay in hospital	94
10. Comparison of effects of treatment with Nigella Sativa 250 mg and 500 mg on objective signs of withdrawal from opioids during 12 days' stay in hospital	97

(Continued.....)

	PAGE
11. Comparison of effects of treatment with Nigella Sativa 250 mg and 500 mg on subjective symptoms of protracted abstinence in patients with opioid dependence during 12 weeks of follow up	103
12. Comparison of effects of treatment with Nigella Sativa 250 mg and 500 mg on objective signs of protracted abstinence in patients with opioid dependence during 12 weeks of follow up	105
