

CONTENTS

| CHAPTER | TITLE | PAGE |
|---------|--|------|
| | Acknowledgments | I |
| | Contents | II |
| | List of Tables | VI |
| | List of Figures | X |
| | List of abbreviations | XIII |
| | Abstract | XIV |
| 1 | INTRODUCTION | 1 |
| 2 | REVIEW OF LITRATURE | 4 |
| 2.1 | Seedling age | 4 |
| 2.2 | Nursery Management | 6 |
| 2.3 | System of Rice Intensification | 9 |
| 2.4 | Importance of zinc application | 10 |
| 2.6 | Factors controlling zinc availability | 15 |
| 2.7 | Importance of boron application | 19 |
| 2.12 | Aerobic rice | 23 |
| 3 | MATERIALS AND METHODS | 26 |
| 3.1 | Site and soil | 26 |
| 3.2 | Soil analysis | 26 |
| 3.3 | Treatment and layout of experiments | 28 |
| 3.4 | Metrological data recorded during rice growing seasons | 31 |
| 3.5 | Calculation of amount of water applied | 33 |
| 3.6 | OBSERVATIONS | 33 |
| 3.6.1 | Phonological parameters | 34 |
| 3.6.2 | Yield attributes | 35 |
| 3.6.3 | Quality parameters | 36 |
| 3.7 | Determination of total boron | 38 |

| | | | |
|---|--------|--|----|
| | 3.8 | Estimation of Zinc contents | 38 |
| | 3.9 | Statistical Analysis | 38 |
| 4 | | RESULTS AND DISCUSSION | 39 |
| | 4.1 | Nursery Parameters | 39 |
| | 4.1.1 | Nursery plant population (m ⁻²) | 39 |
| | 4.1.2 | Characteristics of 10-days older seedlings | 39 |
| | 4.1.3 | Characteristics of 20-days older seedlings | 40 |
| | 4.1.4 | Characteristics of 30-days older seedlings | 43 |
| | 4.1.5 | Characteristics of 40-dyas older seedlings | 45 |
| | 4.2 | DISCUSSION | 48 |
| | 4.3 | Crop Growth Analysis | 50 |
| | 4.3.1 | Leaf Area Index (LAI) (m ⁻²) | 50 |
| | 4.3.2 | Leaf Area Duration (LAD) (days) | 53 |
| | 4.3.3 | Crop Growth Rate (CGR) (days) | 52 |
| | 4.3.4 | Total dry matter accumulation (TDM) (g m ⁻²) | 53 |
| | 4.3.5 | Net Assimilation Rate (NAR) (g m ⁻²) | 53 |
| | 4.4 | DISCUSSION | 62 |
| | 4.5 | AGRONOMIC TRAITS | 64 |
| | 4.5.1 | Total tillers (m ⁻²) | 64 |
| | 4.5.2 | Number of productive tillers (m ⁻²) | 64 |
| | 4.5.3 | Plant Height (cm) | 66 |
| | 4.5.4 | 1000-grain weight | 65 |
| | 4.5.5 | Panicle length (cm) | 66 |
| | 4.5.6 | Kernel/ panicle | 66 |
| | 4.5.7 | Straw yield (t/ha) | 66 |
| | 4.5.8 | Paddy yield (t/ha) | 67 |
| | 4.5.9 | Biological yield (t/ha) | 67 |
| | 4.5.10 | Harvest Index | 68 |
| | 4.5.11 | Nitrogen concentration (grain + straw) g/kg | 68 |

| | | |
|--------|--|-----|
| 4.6 | Economic Return | 81 |
| 4.7 | DISCUSSION | 83 |
| 4.8 | Crop Growth Analysis | 86 |
| 4.8.1 | Final germination count (m ⁻²) | 86 |
| 4.8.2 | Leaf area index (LAI) (m ⁻²) | 86 |
| 4.8.3 | Leaf Area Duration (LAD) (days) | 86 |
| 4.8.4 | Crop Growth Rate (CGR) (days) | 87 |
| 4.8.5 | Total dry matter accumulation (TDM) (g m ⁻²) | 87 |
| 4.8.6 | Net Assimilation Rate (NAR) (g m ⁻²) | 88 |
| 4.8.7 | Chlorophyll contents (mg/g fw) | 97 |
| 4.9 | DISCUSSION | 103 |
| 4.10 | AGRONOMIC TRAITS | 104 |
| 4.10.1 | Plant height (cm) | 104 |
| 4.10.2 | Total tillers (m ⁻²) | 104 |
| 4.10.3 | Productive tillers (m ⁻²) | 105 |
| 4.10.4 | Panicle length (cm) | 105 |
| 4.10.5 | 1000-kernel weight (g) | 106 |
| 4.10.6 | Straw yield (t/ha) | 106 |
| 4.10.7 | Paddy yield (t/ha) | 107 |
| 4.10.8 | Biological yield (t/ha) | 107 |
| 4.10.9 | Harvest Index (%) | 108 |
| 4.11 | Nutrient contents (mg/kg) | 108 |
| 4.12 | Water Productivity (g/liter) | 122 |
| 4.13 | Economic Return | 122 |
| 4.14 | DISCUSSION | 125 |
| 4.15 | QUALITY PARAMETERS | 128 |
| 4.15.1 | Sterility (%) | 128 |
| 4.15.2 | Abortive Kernels (%) | 128 |
| 4.15.3 | Opaque Kernels (%) | 128 |

| | | | |
|---|--------|----------------------|-----|
| | 4.15.4 | Normal Kernels (%) | 130 |
| | 4.15.5 | Amylose contents (%) | 129 |
| | 4.15.6 | Protein contents (%) | 129 |
| | 4.16 | DISCUSSION | 136 |
| 5 | | SUMMARY | 139 |
| | 5.1 | Experiment-1 | 139 |
| | 5.2 | Experiment-2 | 139 |
| | 5.3 | CONCLUSIONS | 142 |
| | 5.4 | FUTURE THRUST | 143 |
| | | LITRATURE CITED | 144 |