

Chapter	Contents	Page No.
I	APPROVAL CERTIFICATE BY SUPERVISORY COMMITTEE	v
II	THESIS RELEASE FORM	vii
III	ACKNOWLEDGEMENTS	viii
IV	LIST OF TABLES	ix
V	LIST OF FIGURES	xi
VI	APPENDICES	xviii
VII	ABSTRACT	xix
I	INTRODUCTION	1
1.2	Taxonomy	3
1.3	Biotype	4
1.4	Life history	4
1.5	Objectives	7
II	REVIEW OF LITERATURE	9
2.1	Biology	9
2.2	Temperature	14
2.3	Starvation	16
2.4	Dispersal and distribution	18
2.5	Leaf age	22
2.6	Life table	25
2.7	Predators	29
2.8	Parasitoids	31
2.9	Host plants	37
III	MATERIALS AND METHODS	45
3.1	Biology of <i>Bemisia tabaci</i> (Genn.)	45
3.1.1	Culture development	45
3.1.2	On melon and cucumber	46
3.2	Fecundity, egg hatching and development of <i>Bemisia tabaci</i> under constant condition	47
3.2.1	In laboratory	47
3.2.2	In field temperature	48
3.3	Starvation capacity of newly hatched whitefly crawlers	49
3.4	Distribution of premature stages of whitefly, <i>Bemisia tabaci</i> on melon leaf.	50
3.5	Effect of leaf maturity on oviposition and nymphal mortality of <i>Bemisia tabaci</i> .	52
3.6.1	Life table of whitefly, <i>Bemisia tabaci</i> in field conditions	55
3.6.2	Preparation of life table	57
3.7	Seasonal abundance of parasitoids on Whitefly <i>Bemisia tabaci</i> at Dadu, Tandojam and Thatta.	60

3.7	Sampling (dispersal) of <i>Bemisia tabaci</i> adults in melon field	62
3.7.1	At different ground levels	62
3.7.2	On melon creeper	62
3.8	Preferential distribution of Whitefly, <i>Bemisia tabaci</i> on cucurbit and non-cucurbit crops	63
3.9.1	Seasonal population variation of whitefly, <i>Bemisia tabaci</i> on various cucurbit crops in different agro-zones of Sindh	65
V	EXPERIMENTAL RESULT	68
4.1	Biology of <i>Bemisia tabaci</i> on cucurbits	68
4.1.1	On cucumber	68
4.1.2	On musk melon	68
4.2.1	Fecundity, egg hatching and development of <i>Bemisia tabaci</i> under constant condition	72
4.2.2	Constant temperatures (Laboratory)	72
4.2.3	Field temperature	74
4.3	Starvation capacity of newly hatched whitefly crawlers	84
4.4	Distribution of premature stages of whitefly, <i>Bemisia tabaci</i> on melon leaf.	87
4.5	Effect of leaf maturity on oviposition and nymphal mortality of <i>Bemisia tabaci</i>	91
4.6	Life table of <i>B. tabaci</i> in field condition	96
4.6.1	Egg mortality	96
4.6.2	Larval Predation and Parasitism	96
4.6.3	Life table 2000	98
4.6.4	Overall mortality and survival 2000	110
4.6.5	Life table 2001	111
4.6.6	Overall mortality and survival 2001	122
4.6.7	Life table 2002	127
4.6.8	Overall mortality and survival 2002	139
4.6.9	Comparative mortality and survival from 2000-2002	142
4.6.10	Over all mortality and survival of various life stages 2000-2002	143
4.7	Seasonal abundance of parasitoids on Whitefly, <i>Bemisia tabaci</i> at Dadu, Tandojam and Thatta	147
4.7.1	Parasitism percentage	152
4.7.2	<i>B. tabaci</i> and parasitoids ratio	155
4.8	Dispersal (sampling) of <i>Bemisia tabaci</i> adults in melon field	157
4.8.1	Indirect Count	157
4.8.2	Whitefly activity at different ground levels	158
4.8.3	Direct Count	159
4.8.4	Suitable timing of <i>B. tabaci</i> sampling	160

4.9	Preferential distribution of Whitefly, <i>Bemisia tabaci</i> on cucurbit and non-cucurbit crops	162
4.10	Seasonal population variation of whitefly, <i>Bemesia tabaci</i> on various cucurbit crops in different agro-zones of Sindh.	165
4.10.1	Population variation during 2000.	165
4.10.1.a	The population of whitefly on musk melon creepers in various seasons	165
a.1	Spring season	165
a.2	Summer season	165
a.3	Autumn season	166
4.10.1.b	Population of whitefly on Indian squash (Tinda) creepers	168
b.1	Spring season	168
b.2	Summer season	169
b.3	Autumn season	169
4.10.1.c	Population of whitefly on water melon creepers	171
c.1	Spring and summer seasons	171
4.10.1.d	Population of whitefly on cucumber creepers	173
d.1	Spring season	173
d.2	Summer season	173
d.3	Autumn season	173
4.10.1.e	The population of whitefly on ridge gourd (Tori) creepers	174
e.1	Spring, summer and autumn seasons	174
4.10.1.f	The population of whitefly on bottle gourd (Kadu) creepers	175
4.10.1.g	Population of whitefly on bitter gourd (karella) creepers	176
g.1	Spring and summer season	176
4.10.2	Population variation in 2001	183
4.10.2.a	Whitefly on cucumber	184
a.1	Spring population	184
a.2	Summer population	184
a.3	Autumn population	185
4.10.2.b	Whitefly population on Indian squash	187
b.1	Spring population	187
b.2	Summer population	187
b.3	Autumn population	188
4.10.2.c	Whitefly population on water melon	190
4.10.2.d	Whitefly population on musk melon	192

d.1	Spring population	192
d.2	Summer population	192
d.3	Autumn population	193
V	DISCUSSOIN	201
5.1	Biology of <i>Bemisia tabaci</i> (Genn.) on musk melon and cucumber	201
5.2.1	Fecundity, egg hatching and development under constant condition	203
5.2.2	In laboratory	203
5.2.3	Field conditions	205
5.3	Starvation capacity	206
5.4	Distribution of premature stages on melon leaf	207
5.5	Effect of leaf maturity on oviposition and nymphal mortality	208
5.6	Life table 2001-2002	209
5.7	Seasonal abundance of parasitoids	217
5.8	Sampling (dispersal) of adults in melon field.	219
5.8.1	At ground levels	219
5.8.2	On melon creeper	221
5.9	Preferential distribution on cucurbits and non-cucurbits	221
5.10	Seasonal population variation of <i>B. tabaci</i> on cucurbit creepers during 2000 and 2001	223
5.10.1	Seasonal population variation in 2000	223
5.10.2	Population variation 2001	226
5.10.3	Comparative population 2000 and 2002	228
VI	SUMMARY, CONCLUSION AND RECOMMEDATIONS	230
6.1	Summary	230
6.2	Conclusions	237
6.3	Recommendations	237
VII	LITERATURE CITED	239
	Appendices	271