

TABLE OF CONTENTS

1– INTRODUCTION	1-78
<hr/>	
1.1. CRUDE OIL	2
1.1.1. Components of crude oil	3
1.1.1.1. Saturates	3
<i>1.1.1.1.1. Alkanes or Paraffins (Type formula C_nH_{2n+2})</i>	3
<i>1.1.1.1.2. Cycloalkanes (Naphthenes)(type formula C_nH_{2n-2})</i>	3
1.1.1.2. Aromatics	3
1.1.1.3. Resins and Asphaltenes	5
1.1.2. Characteristics of crude oil	5
1.1.3. Prevalence in the environment	5
1.1.4. Perilous effects of crude oil	5
1.2. POLYCYCLIC AROMATIC HYDROCARBONS	6
1.2.1. Characteristics of PAHs	7
1.2.2. Prevalence in the environment	10
1.2.2.1. Petrogenic PAHs	10
1.2.2.2. Pyrogenic PAHs	10
1.2.2.3. Biogenic PAHs	10
1.2.3. Perilous effects of PAHs	11
1.2.3.1. Nap	13
1.2.3.2. Phe	13
1.2.3.3. Fla	14
1.3. BIODEGRADATION	14
1.3.1. Anaerobic biodegradation	17
1.3.2. Aerobic biodegradation	18
1.3.2.1. Bacterial degradation of PAHs	19

1.3.2.2. Screening of PAHs degrading bacteria	26
<i>1.3.2.2.1. Oxidation-reduction indicator (redox indicator)</i>	26
<i>1.3.2.2.2. Accumulated coloured metabolites</i>	26
<i>1.3.2.2.3. Spray plate methods</i>	26
<i>1.3.2.2.4. Radioactive labeling of PAHs</i>	26
<i>1.3.2.2.5. Microtiter plate assay</i>	27
1.3.2.3. Bacterial metabolic pathways of PAHs degradation	27
<i>1.3.2.3.1. Nap degradation pathways</i>	27
<i>1.3.2.3.2. Phe degradation pathways</i>	31
<i>1.3.2.3.3. Fla degradation pathways</i>	34
<i>1.3.2.3.4. Other HMW PAHs degradation</i>	38
1.4. AROMATIC RING HYDROXYLATING DIOXYGENASE ENZYMES (ARHDs)	39
1.4.1. Oxidases	39
1.4.2. Oxygenases	40
1.4.2.1. Monooxygenases	40
1.4.2.2. Dioxygenases	40
<i>1.4.2.2.1. Classification of ARHDs</i>	42
<i>1.4.2.2.2. Aromatic ring dioxygenase-expressing bacteria (ARDB)</i>	47
1.5. PAHs CATABOLIC GENES	48
1.5.1. PAHs catabolic genes of Gram-negative bacteria	48
1.5.1.1. <i>nah</i> -like genes of <i>Pseudomonas</i> sp. strains	49
1.5.1.2. <i>phd</i> genes of <i>Comamonas</i> sp. strains	52
1.5.1.3. <i>nag</i> genes of <i>Ralstonia</i> sp. strain	54
1.5.1.4. <i>phn</i> genes of <i>Burkholderia</i> sp. strain	54
1.5.1.5. Genes of <i>Sphingomonas</i> sp. strains	56
1.5.2. PAHs catabolic genes of Gram-positive bacteria	58

1.5.2.1. <i>phd</i> genes of <i>Nocardioides</i> sp. strain	58
1.5.2.2. <i>nid</i> genes of <i>Mycobacterium</i> sp. strain	59
1.5.2.3. <i>dbf/fln</i> genes of <i>Terrabacter</i> sp. strain	62
1.5.2.4. <i>nar</i> genes of <i>Rhodococcus</i> sp. strain	63
1.5.2.5. Genes of <i>Kocuria</i> sp. strains	66
1.6. PCR BASED APPROACH TO CHARACTERIZE PAHs DEGRADING BACTERIAL COMMUNITIES	70
1.6.1. Alpha subunit based primers	71
1.6.2. [Fe ₂ -S ₂] Rieske Center based primers	73
1.7. FACTORS INFLUENCING BIODEGRADATION OF PAHs	74
1.7.1. Temperature	76
1.7.2. pH	76
1.7.3. Nutrients	77
1.7.4. Concentration of target PAHs	78
2- OBJECTIVES	79-80
<hr/>	
3- MATERIALS AND METHODS	81-105
<hr/>	
3.1. Sample collection	82
3.2. Enrichment and isolation	82
3.2.1. Media and chemicals	82
3.2.2. Enrichment on low molecular weight (LMW) PAHs	83
3.2.3. Enrichment on high molecular weight (HMW) PAHs	83
3.3. Colonial and cellular morphology	84
3.4. Screening for growth on PAHs	84
3.4.1. Microtiter plate assay to screen isolated bacterial strains	85

3.4.2. Spray plate method to screen bacterial consortia	86
3.5. Selection of bacterial cultures	86
3.6. Identification	86
3.6.1. Preliminary identification	86
3.6.2. 16S rRNA gene analysis	86
3.7. Maintenance and long term preservation of cultures	87
3.8. Biodegradation of PAHs and crude oil	87
3.8.1. Bacterial strains isolated from Nap enriched samples	87
<i>3.8.1.1. Biodegradation of Nap as sole carbon source</i>	87
<i>3.8.1.2. Biodegradation of Nap with additional carbon source</i>	88
<i>3.8.1.3. Biodegradation of Phe as sole carbon source</i>	88
<i>3.8.1.4. Biodegradation of Phe with additional carbon source</i>	89
<i>3.8.1.5. Biodegradation of Fla as sole carbon source</i>	89
<i>3.8.1.6. Biodegradation of Fla with additional carbon source</i>	90
<i>3.8.1.7. Effect of autoclaving on composition of ILCO</i>	90
<i>3.8.1.8. Growth on ILCO coated minimal agar</i>	91
<i>3.8.1.9. Biodegradation of ILCO as sole carbon source</i>	91
<i>3.8.1.10. Biodegradation of ILCO with additional carbon source</i>	91
3.8.2. Bacterial consortia enriched on Fla	92
<i>3.8.2.1. Fla degradation by consortia enriched on minimal medium (MO)</i>	92
<i>3.8.2.2. Fla degradation by consortia enriched on YE added minimal medium (MM)</i>	93
3.8.2.2.1. Time scale biodegradation of Fla	93
3.8.2.2.2. Role of microeukaryotes and prokaryotes in biodegradation of Fla	94
3.8.2.2.3. Aromatic ring dioxygenase-expressing bacteria (ARDB)	95
<i>3.8.2.3. Effect of long incubation on Fla biodegradation of LPMO consortium</i>	
<i>(MO enriched)</i>	95
3.8.3. Bacterial strain CMGCZ enriched on Fla	96
<i>3.8.3.1. Biodegradation of Fla as sole carbon source</i>	96

3.8.3.2. <i>Biodegradation of Fla with additional carbon source</i>	96
3.8.3.3 <i>Biodegradation of Fla in enriched LB medium</i>	96
3.8.3.4. <i>Biodegradation of Nap as sole carbon source</i>	97
3.8.3.5. <i>Biodegradation of Nap with additional carbon source</i>	97
3.8.3.6. <i>Biodegradation of Phe as sole carbon source</i>	97
3.8.3.7. <i>Biodegradation of Phe with additional carbon source</i>	98
3.8.3.8. <i>Biodegradation of ILCO as sole carbon source</i>	98
3.8.3.9. <i>Biodegradation of ILCO with additional carbon source</i>	98
3.8.4. Optimization of biodegradation of PAHs	99
3.8.4.1. <i>Temperature</i>	99
3.8.4.2. <i>pH</i>	99
3.8.4.3. <i>Concentration of Fla</i>	99
3.9. Genetic studies	100
3.9.1. Isolation of plasmid DNA	100
3.9.1.1. <i>Alkaline lysis method of Birnboim and Doly (1979)</i>	100
3.9.1.2. <i>Qiagen plasmid purification kit</i>	101
3.9.1.3. <i>Agarose gel electrophoresis</i>	101
3.9.2. Isolation of genomic DNA	101
3.9.2.1. <i>Beads beating (physical extraction) method</i>	101
3.9.3. PCR amplification of PAH dioxygenase gene	102
3.9.3.1. <i>PCR with genomic DNA</i>	103
3.9.3.2. <i>PCR with single bacterial colony</i>	104
3.9.3.3. <i>Agarose gel electrophoresis</i>	104
3.9.4. Purification of amplified PCR product	104
3.9.5. Sequencing of purified PCR product	105
3.9.6. Sequence analysis	105

4– RESULTS	106-230
<hr/>	
4.1. Enrichment and isolation	107
4.1.1. Enrichment on low molecular weight (LMW) PAHs	107
4.1.2. Enrichment on high molecular weight (HMW) PAHs	107
4.2. Colonial and cellular morphology	110
4.3. Screening for growth on PAHs	114
4.3.1. Microtiter plate assay to screen isolated bacterial strains	114
4.3.2. Spray plate method to screen bacterial consortia	117
4.4. Selection of bacterial cultures	121
4.5. Identification	121
4.5.1. Preliminary identification	121
4.5.2. 16S rRNA identification	123
4.6. Biodegradation of PAHs and crude oil	127
4.6.1. Bacterial strains isolated from Nap enriched samples	127
4.6.1.1. <i>Biodegradation of Nap as sole carbon source</i>	127
4.6.1.2. <i>Biodegradation of Nap with additional carbon source</i>	127
4.6.1.3. <i>Biodegradation of Phe as sole carbon source</i>	132
4.6.1.4. <i>Biodegradation of Phe with additional carbon source</i>	132
4.6.1.5. <i>Biodegradation of Fla as sole carbon source</i>	135
4.6.1.6. <i>Biodegradation of Fla with additional carbon source</i>	135
4.6.1.7. <i>Effect of autoclaving on composition of ILCO</i>	138
4.6.1.8. <i>Growth on ILCO coated minimal agar</i>	141
4.6.1.9. <i>Biodegradation of ILCO as sole carbon source</i>	141
4.6.1.10. <i>Biodegradation of ILCO with additional carbon source</i>	141
4.6.2. Bacterial consortia enriched on Fla	143
4.6.2.1. <i>Consortia enriched on MO</i>	143
4.6.2.1.1. <i>Biodegradation of Fla</i>	143

4.6.2.2. Consortia enriched on MM	143
4.6.2.2.1 Biodegradation of Fla	143
4.6.2.2.2. Time scale biodegradation of Fla	151
4.6.2.2.3. Role of microeukaryotes in biodegradation of Fla	156
4.6.2.2.4. Role of prokaryotes in biodegradation of Fla	160
4.6.2.2.5. Aromatic ring dioxygenase-expressing bacteria (ARDB)	164
4.6.2.2.5.1. ARDB of LPMM consortium	164
4.6.2.2.5.2. ARDB of LPMO consortium	164
4.6.2.2.5.3. ARDB of long incubated LPMO consortium	164
4.6.2.2.5.4. ARDB of HPMM consortium	169
4.6.2.2.5.5. ARDB of HPMO consortium	169
4.6.2.3. Effect of long incubation on Fla biodegradation of LPMO consortium	171
4.6.3. Bacterial strain <i>R. erythropolis</i> CMGCZ enriched on Fla	176
4.6.3.1. Biodegradation of Fla as sole carbon source	176
4.6.3.2. Biodegradation of Fla with additional carbon source	179
4.6.3.3. Biodegradation of Fla in LB medium	182
4.6.3.4. Biodegradation of Nap as sole carbon source	185
4.6.3.5. Biodegradation of Nap with additional carbon source	185
4.6.3.6. Biodegradation of Phe as sole carbon source	188
4.6.3.7. Biodegradation of Phe with additional carbon source	188
4.6.3.8. Biodegradation of ILCO as sole carbon source	191
4.6.3.9. Biodegradation of ILCO with additional carbon source	191
4.6.4. Optimization of biodegradation of PAHs	195
4.6.4.1. Temperature	195
4.6.4.2. pH	200
4.6.4.3. Concentration of Fla	205
4.7. Genetic studies	213
4.7.1. Isolation of plasmid DNA	213

4.7.2. Isolation of genomic DNA	214
4.7.3. PCR amplification of PAH dioxygenase gene	214
<i>4.7.3.1. PCR with genomic DNA</i>	214
<i>4.7.3.2. PCR with single bacterial colony</i>	217
4.7.4. Purification of amplified PCR product	224
4.7.5. Sequencing of purified PCR product	225
4.7.6. Sequence analysis	225
5- DISCUSSION	231-257
<hr/>	
6- CONSLUSION & FINDINGS	258-260
<hr/>	
7- FUTURE STANCES	261-262
<hr/>	
8- REFERENCES	263-318
<hr/>	
9- APPENDICE	319-325
<hr/>	